



(CASE REPORT)



Combination of ketamine, corticosteroids and sevoflurane inhibits the risk of bronchospasm in intubated children under general anesthesia

Maria I Dalamagka *

Department of Anesthesia, General Hospital of Larisa, Greece.

GSC Advanced Research and Reviews, 2022, 11(01), 165–166

Publication history: Received on 09 March 2022; revised on 17 April 2022; accepted on 17 April 2022

Article DOI: <https://doi.org/10.30574/gscarr.2022.11.1.0108>

Abstract

Asthma in children is associated with significant morbidity. Children with severe asthma are at increased risk for adverse outcomes including medication-related side effects, life-threatening exacerbations, and impaired quality of life. In the study, an asthmatic child with a recent cold, received general anesthesia for emergency surgery centered on sevoflurane, corticosteroids and ketamine. The purpose of this study is to demonstrate the beneficial effects of the combination of sevoflurane, ketamine and corticosteroids in asthmatic children and to prevent complications when they are given general anesthesia for emergency surgery.

Keywords: Corticosteroids; Sevoflurane; Bronchospasm; Ketamine; Anesthesia; Astma

1. Introduction

Asthma is a chronic respiratory disease that affects people of all ages and is characterized by episodic and reversible attacks of wheezing, chest tightness, shortness of breath, and coughing. According to the ATS/ERS guideline, severe asthma is defined as asthma which requires treatment with high dose inhaled corticosteroids (ICS) plus a second controller (and/or systemic corticosteroid) to prevent it from becoming “uncontrolled” or remains “uncontrolled” despite this therapy [1-7].

2. Case Study

A 6-year-old child weighing 30 kg with a recent cold, nasal congestion and hearing of both amphibians and with a known history of frequent asthma attacks underwent emergency appendectomy. It was given intravenously before the introduction of anesthesia solu medrol 60 mg, Dexaton 3 mg, onda 3 mg. Introduction to anesthesia was by intravenous administration of Fentanyl 60 mcg, Propofol 120 mg, Esmeron 30 mg. After intubation, intrabronchial aspiration was performed, intrabronchial Flixotide 50 mcg και Aerolin 100 mcg was given and mechanically ventilated with 50% N₂O and 2% sevoflurane. Ketamine 6 mg was given intravenously, plus 30 mcg Fentanyl, Apotel 350 mg, and morphine 1.5 mg. The monitoring included ECG, NBP, SpO₂ and the ventilation model in Drager machine was Volune Control. Towards the end of the operation and with pure inhaled oxygen he was put in a Pressure Control model and the awakening was done smoothly after intravenous Bridium 0.6 mg.

3. Management and Outcome

The common denominator underlined in all forms of asthma is bronchial hyperresponsiveness to various stimuli. Inhaled glucocorticoids have long been used as a first-line treatment for persistent pediatric asthma, as they are the

* Corresponding author: Maria I Dalamagka
Anesthesia Department, General Hospital of Larisa, Greece.

most effective intervention for the treatment of asthma. Thus, Solu medrol (kg x 2), Dexaton (up to 0.1x kg) and Flixotide 50 mcg and Aerolin 100 mcg were administered intravenously. Ketamine also causes bronchodilation and was administered at its appropriate titrated dose (0.2 x kg), as anesthesia was maintained with sevoflurane which does not irritate the respiratory system.

4. Discussion

Corticosteroids have inhibitory properties in many effects on many stem cells and inflammatory cells, which are activated in asthma. Inhaled steroids reduce the number and activation of inflammatory cells in the epithelium and submucosa by clogging the damaged epithelium and, potentially, inhibiting the production of proinflammatory cytokines, and reducing the survival time of the epithelium. This action of corticosteroids in combination with the ketamine which has a bronchodilator effect and the property of sevoflurane as it does not irritate the airway, they eliminate bronchospasm and prevent possible laryngospasm.

5. Conclusion

In conclusion, the combination of Ketamine, corticosteroids and sevoflurane has been shown to inhibit possible complications such as bronchospasm when asthmatic children implied in general anesthesia.

Compliance with ethical standards

Acknowledgments

General Hospital of Larisa.

Statement of informed consent

Written informed consent was obtained from the patient for publication of this case report. A copy of the written consent is available.

References

- [1] KF Chung, SE Wenzel, JL Brozek, A Bush, M Castro, PJ Sterk, et al. International ERS/ATS guidelines on definition, evaluation and treatment of severe asthma Eur Respir J. 2014; 43; 343-373.
- [2] HS Zahran, CM Bailey, SA Damon, PL Garbe, PN Breysse. Vital signs: asthma in children – United States, 2001-2016 MMWR Morb Mortal Wkly Rep. 2018; 67: 149-155.
- [3] WG Teague, BR Phillips, JV Fahy, SE Wenzel, AM Fitzpatrick, WC Moore. et al. Baseline features of the severe asthma Research program (SARP III) cohort: differences with age J Allergy Clin Immunol Pract. 2018; 6: 545-554.
- [4] F Rabe, P Nair, G Brusselle, JF Maspero, M Castro, L. Sher et al. Efficacy and safety of dupilumab in glucocorticoid-dependent severe asthma N Engl J Med. 2018; 378: 2475-2485.
- [5] S Gonem, R Berair, A Singapuri, R Hartley, MFM Laurencin, G Bacher, et al. Fevipiprant, a prostaglandin D2 receptor 2 antagonist, in patients with persistent eosinophilic asthma: a single-centre, randomised, double-blind, parallel-group, placebo-controlled trial Lancet Respir Med. 2016; 4: 699-707.
- [6] H Farne, DJ Jackson, SL Johnston Are emerging PGD2 antagonists a promising therapy class for treating asthma? Expert Opin Emerg Drugs. 2016; 21: 359-364.
- [7] J Corren, JR Parnes, L Wang, M Mo, SL Roseti, JM Griffiths, et al. Tezepelumab in adults with uncontrolled asthma N Engl J Med. 2017; 377: 936-946.